

### Characteristics of major non-exploited brown coal deposits

	Name of deposit	Thickness of coal seams (in m)	Depth to the base (m)	Overburden to deposit thickness ratio	Calorific value (kcal/kg)	Ash content (%)	Mean. content of sulfur (%)
1.	Babina - Żarki	10.7	140.0	-	2,229	18.28	1.10
2.	Cybinka	16.6	94.0	5.6	2,236	18.40	1.41
3.	Gubin	10.9	83.7	7.4	2,24	12.86	1.42
4.	Głowaczów	4.8	37.1	6.5	1,82	28.56	0.42
5.	Legnica p. Północ	22.0	207.2	8.7	2,194	18.72	1.42
6.	Legnica p. Wschód	19.8	136.3	7.3	2,206	19.05	1.33
7.	Legnica p. Zachód	21.0	158.8	6.6	2,371	20.10	0.76
8.	Mosty	9.3	105.0	8.6	2,219	17.19	1.63
9.	Piaski	6.1	48.5	7.3	2,082	24.80	1.44
10.	Rogoźno	35.0	195.0	4.6	2,319	18.90	3.99
11.	Rzepin	12.2	97.3	7.9	2,164	15.14	1.20
12.	Sądów	12.2	127.5	10.2	2,196	18.80	1.38
13.	Ścinawa	20.2	207.1	8.9	2,276	12.88	0.48
14.	Torzym	21.4	180.8	7.9	2,27	16.80	1.81
15.	Trzcianka	6.0	80.0	8.6	1,888	28.2	1.84
16.	Złoczew	46.2	259.1	4.5	2,021	21.67	1.18