

Table 2

Characteristics of major non-exploited brown coal deposits

	Name of deposit	Thickness of coal seams (in m)	Depth to the base	Overburden-to deposit thickness ration	Calorific value (kcal/kg)	Ash content (%)	Mean. content of sulfur (%)
1	Babina - Żarki	10.7	140.0	-	2 229	18.28	1.10
2	Cybinka	16.6	94.0	5.6	2 236	18.40	1.41
3	Gubin	10.9	83.7	7.4	2 240	12.86	1.42
4	Głowaczów	4.8	37.1	6.5	1 820	28.56	0.42
5	Legnica p. Północ	22.0	207.2	8.7	2 194	18.72	1.42
6	Legnica p. Wschód	19.8	136.3	7.3	2 206	19.05	1.33
7	Legnica p. Zachód	21.0	158.8	6.6	2 371	20.10	0.76
8	Mosty	9.3	105.0	8.6	2 219	17.19	1.63
9	Piaski	6.1	48.5	7.3	2 082	24.80	1.44
10	Rogóżno	18,7	104,5	4,9	2 244	18,90	3,99
11	Rzepin	12.2	97.3	7.9	2 164	15.14	1.20
12	Sądów	12.2	127.5	10.2	2 196	18.80	1.38
13	Ścinawa	20.2	207.1	8.9	2 276	12.88	0.48
14	Torzym	21.4	180.8	7.9	2 270	16.80	1.81
15	Trzcianka	6.0	80.0	8.6	1 888	28.20	1.84
16	Złoczew	46.2	259.1	4.5	2 021	21.67	1.18