

Table 2

## Characteristics of major non-exploited lignite deposits

	Name of deposit	Thickness of coal seams (in m)	Depth to the base	Overburden-to deposit thickness ratio	Calorific value (kcal/kg)	Ash content (%)	Mean. Content of sulfur (%)
1	Babina - Żarki	10.7	140.0	-	2 229	18.28	1.10
2	Cybinka	8.8	78.0	7.9	2 251	17.16	1.32
3	Gubin	7.6	99.7	9.3-16.0	2 240	12.86	1.42
4	Gubin 1	10.4	85.3	9.0	2 183	20.37	1.95
5	Gubin 2	13.3	127.6	8.2-8.7	2 246	16.86	2.98
6	Głowaczów	4.8	37.1	6.5	1 820	28.56	0.42
7	Legnica-pole Północne	22.0	207.2	8.7	2 194	18.72	1.42
8	Legnica-pole Wschodnie	18.1	155.4	7.6	2 090	21.80	0.42
9	Legnica-pole Zachodnie	21.0	158.8	6.6	2 254	18.14	0.74
10	Mosty	9.1	82.7	8.0	2 219	17.19	1.63
11	Piaski	6.1	48.5	7.3	2 072	24.90	1.44
12	Rogóźno	18,7	104,5	4.9	2 241	24.54	3.79
13	Rzepin	12.2	97.3	7.9	2 164	15.14	1.20
14	Sądów	12.2	127.5	10.2	2 196	19.16	1.39
15	Ścinawa	20.2	207.1	8.9	2 275	12.88	0.64
16	Torzym	21.4	180.8	7.9	2 270	16.80	1.81
17	Trzcianka	4.2	45.9	11.9	2 004	23.07	1.84
18	Złoczew	51.4	266.6	5.0	1 968	21.29	2.14